

Article

Induction of oxidative stress causes functional alterations in mouse urothelium via a TRPM8-mediated mechanism: implications for aging

Nocchi, Linda, Daly, Donna, Chapple, Christopher and Grundy, David

Available at <http://clock.uclan.ac.uk/17868/>

Nocchi, Linda, Daly, Donna, Chapple, Christopher and Grundy, David (2014) Induction of oxidative stress causes functional alterations in mouse urothelium via a TRPM8-mediated mechanism: implications for aging. Aging Cell, 13 (3). pp. 540-550. ISSN 1474-9718

It is advisable to refer to the publisher's version if you intend to cite from the work.

<http://dx.doi.org/10.1111/accel.12208>

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>



Induction of oxidative stress causes functional alterations in mouse urothelium via a TRPM8-mediated mechanism: implications for aging

Linda Nocchi,

1

Donna M. Daly,

1

Christopher Chapple

2

and

David Grundy

1

1

Department of Biomedical Science, University of Sheffield, Western Bank, Sheffield, S10 2TN, UK

2

Department of Urology, Royal Hallamshire Hospital, Glossop Road, Sheffield, S10 2JF, UK

Summary

The incidence of bladder conditions such as overactive bladder syndrome and its associated urinary incontinence is highly prevalent in the elderly. However, the mechanisms underlying these disorders are unclear. Studies suggest that the urothelium forms a 'sensory network' with the underlying innervation, alterations in which, could compromise bladder function. As the accumulation of reactive oxygen species can cause functional alterations with age, the aim of this study was to investigate whether oxidative stress alters urothelial sensory signalling and whether the mechanism underlying the effect of oxidative stress on the urothelium plays a role in aging. Five-month-old (young) and 24-month-old (aged) mice were used. H

2

0

2

, used to induce

oxidative stress, resulted in an increase in bladder afferent nerve activity and urothelial intracellular calcium in preparations from young mice. These functional changes were concurrent with upregulation of TRPM8 in the urothelium. Moreover, application of a TRPM8 antagonist significantly attenuated the H

2

0

2

-induced

calcium responses. Interestingly, an upregulation of TRPM8 was also found in the urothelium from aged mice, where high oxidative stress levels were observed, together with a greater calcium response to the TRPM8 agonist WS12. Furthermore, these calcium responses were attenuated by pretreatment with the antioxidant N-acetyl-cysteine. This study shows that oxidative stress affects urothelial function involving a TRPM8-mediated